

# Trauma and psychotic experiences: transnational data from the World Mental Health Survey

John J. McGrath, Sukanta Saha, Carmen C. W. Lim, Sergio Aguilar-Gaxiola, Jordi Alonso, Laura H. Andrade, Evelyn J. Bromet, Ronny Bruffaerts, José M. Caldas de Almeida, Graça Cardoso, Giovanni de Girolamo, John Fayyad, Silvia Florescu, Oye Gureje, Josep M. Haro, Norito Kawakami, Karestan C. Koenen, Viviane Kovess-Masfety, Sing Lee, Jean-Pierre Lepine, Katie A. McLaughlin, Maria E. Medina-Mora, Fernando Navarro-Mateu, Akin Ojagbemi, Jose Posada-Villa, Nancy Sampson, Kate M. Scott, Hisateru Tachimori, Margreet ten Have, Kenneth S. Kendler, Ronald C. Kessler, on behalf of the WHO World Mental Health Survey Collaborators\*

## Background

Traumatic events are associated with increased risk of psychotic experiences, but it is unclear whether this association is explained by mental disorders prior to psychotic experience onset.

## Aims

To investigate the associations between traumatic events and subsequent psychotic experience onset after adjusting for post-traumatic stress disorder and other mental disorders.

## Method

We assessed 29 traumatic event types and psychotic experiences from the World Mental Health surveys and examined the associations of traumatic events with subsequent psychotic experience onset with and without adjustments for mental disorders.

## Results

Respondents with any traumatic events had three times the odds of other respondents of subsequently developing

psychotic experiences (OR=3.1, 95% CI 2.7–3.7), with variability in strength of association across traumatic event types. These associations persisted after adjustment for mental disorders.

## Conclusions

Exposure to traumatic events predicts subsequent onset of psychotic experiences even after adjusting for comorbid mental disorders.

## Declaration of interest

In the past 3 years, R.C.K. received support for his epidemiological studies from Sanofi Aventis, was a consultant for Johnson & Johnson Wellness and Prevention, and served on an advisory board for the Johnson & Johnson Services Inc Lake Nona Life Project. He is a co-owner of DataStat, Inc, a market research firm that carries out healthcare research.

## Copyright and usage

© The Royal College of Psychiatrists 2017.

Extensive evidence indicates that exposure to childhood trauma or other adversities is associated with increased risk of subsequent psychotic experiences.<sup>1</sup> In a recent analysis of data from the World Health Organization (WHO) World Mental Health (WMH) surveys, McGrath *et al*<sup>2</sup> found that childhood adversities, such as sexual or physical abuse, were associated with increased risks of subsequent psychotic experiences even after adjustment for lifetime comorbid mental disorders. Exposure to traumatic events later in life has also been linked to increased risk of subsequent psychotic experiences. Cross-sectional<sup>3</sup> and longitudinal<sup>4,5</sup> studies have found especially high psychotic experience risk associated with traumatic events involving interpersonal violence.<sup>6,7</sup> There is also evidence of a dose–response relationship between number of traumatic events and risk of subsequent psychotic experiences.<sup>6,8</sup> However, many traumatic events are predicted by prior mental disorders and associated with an increased risk of subsequent mental disorders, most notably post-traumatic stress disorder (PTSD).<sup>9–11</sup> We recently demonstrated that the associations of psychotic experiences with mental disorders (including PTSD) are often bidirectional.<sup>12</sup> Thus, it seems reasonable to hypothesise that the associations of traumatic events with subsequent onset of psychotic experiences might be influenced by comorbid mental disorders.<sup>13</sup> This could be true either because lifetime mental disorders with onsets prior to traumatic events are associated both with increased risk of

subsequent traumatic event exposure (such as bipolar disorder associated with increased risk of interpersonal violence, attention-deficit hyperactivity disorder (ADHD) associated with increased risk of motor vehicle collisions) and because traumatic events are associated with increased risk of subsequent mental disorders (both PTSD and other disorders). The main aim of this study was to examine the associations of type and number of traumatic events with the subsequent onset of psychotic experiences across 16 WMH countries and to evaluate the extent to which these associations were explained statistically by PTSD and other mental disorders.

## Method

### Samples

The WMH surveys are a coordinated set of community epidemiological surveys administered in probability samples of the non-institutionalised civilian household population in countries throughout the world ([www.hcp.med.harvard.edu/WMH](http://www.hcp.med.harvard.edu/WMH)).<sup>14</sup> We examined data from 16 WMH surveys that included both a psychosis module and items related to trauma exposure. These 16 countries are distributed across North and South America (Colombia, Mexico, Peru, Brazil, USA); Africa (Nigeria); the Middle East (Lebanon); the South Pacific (New Zealand) and Europe (Belgium, France, Germany, Italy, The Netherlands, Portugal, Romania, Spain). The majority of these surveys were based on multistage, clustered area probability household

\*See Appendix for list of collaborators.

sampling designs, the exceptions being Belgium, Germany and Italy, which used municipal resident registries to select respondents (online Table DS1). The weighted (by sample size) average response rate across all 16 surveys was 70.5%.

In keeping with previous studies of psychotic experiences,<sup>2,6,12</sup> we made the *a priori* decision to exclude individuals reporting psychotic experiences who screened positive for possible schizophrenia/psychosis and manic depression/mania (i.e. respondents who were either told by a doctor that their psychotic experiences were caused by these conditions or who were treated with antipsychotic medications for these symptoms). This resulted in excluding 130 respondents (0.5% of all respondents), leaving 24 464 respondents for this study (online Table DS1).

## Procedures

All WMH interviews were conducted face-to-face by trained lay interviewers in the homes of respondents. Informed consent was obtained before beginning the interview. Procedures for obtaining informed consent and data protection (ethical approvals) were reviewed and approved by the institutional review boards of the collaborating organisations in each country.<sup>15</sup> Standardised interviewer training and quality control procedures were used consistently in the surveys. All WMH interviews had two parts. Part I, administered to all respondents, contained assessments related to core mental disorders (depression, mania, panic disorder, social phobia, specific phobia, agoraphobia, generalised anxiety disorder, substance use). Part II, which included other mental disorders, traumatic events and psychotic experiences, was administered to respondents who met lifetime criteria for any part I disorder and a random proportion of the rest. Respondents in the part II sample were weighted by the inverse of their probability of selection to restore representativeness. Additional weights were used to adjust for differential probabilities of selection within households, non-response and to match the samples to population sociodemographic distributions.<sup>15</sup>

## Data collection and data items

The instrument used in the WMH surveys was the WHO Composite International Diagnostic Interview (CIDI),<sup>15</sup> a validated fully structured diagnostic interview (<https://www.hcp.med.harvard.edu/wmh/cidi/download-the-who-wmh-cidi-instruments/>) designed to assess the prevalence and correlates of a wide range of mental disorders according to the definitions and criteria of both the DSM-IV and ICD-10 diagnostic systems.<sup>16,17</sup> DSM-IV criteria are used in the current report. WHO translation, back-translation and harmonisation protocols were used to adapt the CIDI for use in each participating country.

## Psychotic experiences

The psychosis module included questions about six psychotic experience types – two related to hallucinatory experiences and four related to delusional experiences (online Tables DS2, DS3). The respondents were asked if they ever experienced each psychotic experience (for example ‘Have you ever seen something that wasn’t there that other people could not see?’; ‘Have you ever heard any voices that other people said did not exist?’ etc.). Only psychotic experiences occurring when the person was ‘not dreaming, not half-asleep, or not under the influence of alcohol or drugs’ were included. For those with psychotic experiences, age at onset of psychotic experiences was also assessed.

## Mental disorders

The WMH version of the CIDI assessed lifetime history of 21 mental disorders including mood disorders, anxiety disorders,

behaviour disorders, eating disorder, and substance use disorders (online Table DS4). Clinical reappraisal studies indicate that lifetime diagnoses based on the CIDI have good concordance with diagnoses based on masked clinical interviews.<sup>18</sup>

## Traumatic experiences

The CIDI assessed 29 traumatic events that have been broadly classified into six categories. Details of these categories have been published elsewhere.<sup>19</sup> These categories are: (a) collective violence (civilian in a war zone, refugee, civilian in region of terror, kidnapped, and relief worker in war zone); (b) caused/witnessed bodily harm (purposely injured, tortured, or killed someone, combat experience, accidentally caused serious injury or death, saw atrocities, and witnessed death/dead body or saw someone seriously hurt); (c) interpersonal violence (beaten up by caregiver, witnessed physical fight at home, and beaten up by someone else); (d) intimate partner/sexual violence (raped, sexually assaulted, beaten up by spouse/romantic partner, stalked, traumatic event to loved one, private event, and some other event); (e) accidents/injuries (child with serious illness, natural disaster, life-threatening illness, toxic chemical exposure, other life-threatening accident, and automobile accident); and (f) other traumas (unexpected death of a loved one, mugged or threatened with a weapon, and man-made disaster). Information on age at first exposure associated with each traumatic event was also collected. Of note, this study extends our previous report on the subset of traumatic events that occurred in childhood<sup>2</sup> by evaluating the impact of traumatic events occurring across the lifespan.

## Statistical analysis

Lifetime prevalence of each traumatic event was calculated using cross-tabulation. Discrete-time survival analysis with person-year as the unit of analysis was used to investigate the associations of exposure to traumatic events with the subsequent first onset of psychotic experiences. A person-year data-set was created such that each year in the life of each respondent (up to and including the age at onset of psychotic experience or their age at interview, whichever came first) was treated as a separate observational record. In all analyses the year of psychotic experience onset was coded as 1 and earlier years coded as 0 on a dichotomous outcome variable. Traumatic events were coded as 1 only beginning the year after the onset of traumatic events to ensure that we were not including onsets of traumatic event and psychotic experience occurring in the same year or traumatic events occurring after psychotic experience onset. We estimated 29 bivariate models (i.e. considering only 1 traumatic event type out of 29 types at a time) in predicting psychotic experiences adjusting for age-cohort, gender, person-year dummy variables, and country. We then estimated a series of multivariate models: (a) model 1 included all traumatic events simultaneously as predictors without considering the number of traumatic events (type model); (b) model 2 included number of traumatic events coded as dummy variables (exactly 1, exactly 2, . . . , 5 or more traumatic events) without any information on type of traumatic events (number model); (c) model 3 included both type and number of traumatic events (simple interactive model); and (d) model 4 included type and number of traumatic events along with a separate interaction term between each type and each number of other co-occurring traumatic events (complex interactive model). The latter models showed the ‘marginal effects’ of each type of traumatic event, where a marginal effect is the ‘partial derivative effect’ (a unique contribution) of a specific traumatic event to psychotic experiences while the contribution from all other traumatic events remains constant. Owing to the co-occurring nature of traumatic

events, it is difficult to fractionate the impact of individual traumatic events on psychotic experiences using models 1–3. While model 3 assumed simple interactions, model 4 allowed each type of traumatic event to vary in slope as a function of number of co-occurring traumatic events. The model with the best fit to the data according to the Akaike information criterion (AIC) was carried forward into subsequent examination of the extent to which associations of traumatic events with subsequent psychotic experiences changed when we adjusted for history of PTSD (model 5) and also other temporally prior (to the psychotic experience) mental disorders (model 6).

As the WMH data are both clustered and weighted, the design-based Taylor series linearisation implemented in SUDAAN software was used to estimate standard errors and evaluate the statistical significance of coefficients. Survival coefficients were exponentiated and reported as odds ratios. All significance tests were evaluated using 0.05-level two-sided tests.

## Results

### Prevalence of traumatic events in those with and without psychotic experiences

Of the 24 464 respondents, more than two-thirds ( $n=18\,535$ , 71.8%) reported exposure to at least one traumatic event in their lifetime (online Table DS5). The lifetime prevalence of traumatic event exposure among respondents with psychotic experiences was 90.4% compared with 70.5% in those with no psychotic experiences. Among respondents with psychotic experiences, there was a wide variation in the frequency of exposure to specific types of traumatic events, with the highest prevalence reported for unexpected death of a loved one (45.6%) followed by witnessed death or saw someone seriously hurt (41.8%), and mugged or threatened with weapon (29.9%). In this sample, more than three-quarters (75.8%) of the traumatic events occurred before the onset of psychotic experiences, whereas only 19.2% occurred after psychotic experience onset ( $\chi^2_1=337.3$ ,  $P<0.001$ ) (online Table DS5).

### The associations of type and number of traumatic events with psychotic experiences

Tables 1 and 2 summarise the bivariate and multivariable associations of type and number of traumatic events with subsequent first onset of psychotic experiences. Overall, respondents with any traumatic event had threefold increased odds of subsequent onset of psychotic experiences (OR = 3.1, 95% CI 2.7–3.7) compared with respondents with no traumatic events. In the bivariate models, all but three traumatic event types (civilian in war zone or region of terror, and relief worker in war zone) were significantly associated with elevated odds of subsequent psychotic experiences. The significant odds ratios ranged between 1.6 and 3.5 with the highest being associated with rape (OR = 3.5, 95% CI 2.7–4.4), and the lowest with unexpected death of a loved one (OR = 1.6, 95% CI 1.4–1.9) and natural disaster (OR = 1.6, 95% CI 1.2–2.1).

In model 1, which adjusted for all traumatic events types, 13 of the 29 traumatic event types were significantly associated with onset of psychotic experiences, with odds ratios ranging from 0.3 (95% CI = 0.1–0.8) for relief worker in a war zone to 2.0 (95% CI = 1.0–3.8) for being a refugee. As a set, traumatic events were significantly associated with psychotic experiences ( $\chi^2_{29}=745.7$ ,  $P<0.001$ ), and there was significant variation in the magnitude of associations across traumatic events ( $\chi^2_{28}=71.7$ ,  $P<0.001$ ). With respect to the number of traumatic events (Table 2, model 2), a dose–response relationship between traumatic events and psychotic experiences, was evident with

the odds ratios increasing monotonically as exposure to traumatic events increased ( $\chi^2_5=355.9$ ,  $P<0.001$ ).

In model 3, which adjusted for both type and number of traumatic events, 12 traumatic events were significantly associated with psychotic experiences: 10 with increased odds of subsequent psychotic experiences whereas two traumatic events (‘relief worker in war zone’, ‘civilian in war zone’) were protective with respect to subsequent psychotic experience onset. The odds ratio associated with number of traumatic events were consistently elevated and in the range 1.7–1.3 (Table 2, model 3), indicating that the relative odds of subsequently having psychotic experiences among respondents with more than one prior traumatic event were between 1.7 and 1.3 times the relative odds implied by multiplying together the marginal odds ratios. For example, a respondent who had been exposed to beaten by caregiver, raped, sexually assaulted, and an automobile accident would have an expected odds ratio of subsequent psychotic experiences of 5.0 compared with someone with no traumatic event exposure (i.e.  $1.6 \times 1.6 \times 1.5 \times 1.3$ ). Given that the number of possible traumatic event interaction combinations ( $2^{29}-30=536\,870\,911$ ) greatly exceeded the number of observations, model 3 makes the simplifying assumption that the odds ratios describing interactions are identical for all co-occurring traumatic events involving the same number of traumatic events. However, we found that the test for variation in odds ratios was significant ( $\chi^2_{28}=64.6$ ,  $P<0.001$ ), indicating variation across traumatic event types in the interaction of traumatic event type and number of other traumatic events in predicting psychotic experiences. Thus, we next evaluated a more complex interactive model (model 4) that considered distinct interactions for specific traumatic events together with number of other co-occurring traumatic events.

Table 3 presents a set of complex interactive models with both type and number of traumatic events (model 4) as well as models adjusted for mental disorders (models 5 and 6). Twelve traumatic events were significantly associated with psychotic experience onset in the ‘marginal effects’ models. Interestingly, several traumatic events showed stronger associations with psychotic experience in this model. For example, those who endorsed ‘raped’ had about fivefold increased odds of subsequent onset of psychotic experiences (OR = 4.9, 95% CI 2.5–9.5). The ‘marginal effect’ for number of traumatic events was significant ( $\chi^2_1=5.1$ ,  $P=0.024$ ). The result shows that for every one-unit increment of number of traumatic events, we expect a 10% increase in the odds of subsequent onset of psychotic experiences regardless of the type of traumatic events (OR = 1.1, 95% CI 1.0–1.3).

When we adjusted for PTSD (model 5), the pattern of findings (effect size, direction, and significance) remained remarkably similar. When we adjusted for all mental disorders assessed in the surveys (model 6, including PTSD), 11 of the 29 traumatic events were significantly associated with psychotic experiences. The patterns were generally consistent although for a handful of traumatic events, the odds ratios were slightly attenuated. Overall, these results indicate that the significant associations of specific traumatic events with onset of psychotic experiences were not explained by prior (to the psychotic experiences) mental disorders.

## Discussion

### Main findings

Based on a large data-set from 16 countries, we confirmed that traumatic events are associated with a threefold increased odds of subsequent first onset of psychotic experiences. There was a dose–response relationship between higher numbers of traumatic event types and odds of psychotic experiences. The findings are



<b>Table 1</b> Bivariate and multivariate associations of temporally prior traumatic experiences with subsequent onset of psychotic experiences			
Type of traumatic experiences	Bivariate models <sup>a</sup> OR (95% CI)	M1 – Multivariate type model <sup>b</sup> OR (95% CI)	M3 – Multivariate simple interactive model <sup>c</sup> OR (95% CI)
Collective violence			
Civilian in war zone	1.0 (0.6–1.5)	0.7 (0.4–1.0)	0.6* (0.4–1.0)
Refugee	2.0* (1.1–3.6)	2.0* (1.0–3.8)	1.8 (0.9–3.5)
Civilian in region of terror	1.6 (1.0–2.6)	1.2 (0.7–2.0)	1.1 (0.7–1.9)
Kidnapped	2.3* (1.5–3.7)	1.0 (0.6–1.6)	1.0 (0.6–1.6)
Relief worker in war zone	0.6 (0.3–1.6)	0.3* (0.1–0.8)	0.3* (0.1–0.9)
Caused/witnessed bodily harm			
Purposely injured, tortured or killed someone	2.7* (1.3–5.4)	1.1 (0.5–2.4)	1.1 (0.5–2.5)
Combat experience	1.8* (1.1–3.0)	1.3 (0.8–2.2)	1.3 (0.8–2.1)
Accidentally caused serious injury or death	3.1* (2.2–4.5)	1.6* (1.1–2.3)	1.6* (1.1–2.3)
Saw atrocities	1.7* (1.2–2.4)	0.9 (0.6–1.3)	0.9 (0.7–1.4)
Witnessed death/dead body or saw someone seriously hurt	2.0* (1.8–2.4)	1.4* (1.2–1.7)	1.3* (1.1–1.6)
Interpersonal violence			
Beaten up by caregiver	2.6* (2.2–3.1)	1.7* (1.4–2.1)	1.6* (1.3–1.9)
Witnessed physical fight at home	2.1* (1.8–2.4)	1.2* (1.0–1.5)	1.1 (1.0–1.4)
Beaten up by someone else	2.1* (1.6–2.7)	1.0 (0.8–1.3)	1.0 (0.8–1.4)
Intimate partner/sexual violence			
Raped	3.5* (2.7–4.4)	1.7* (1.2–2.3)	1.6* (1.2–2.3)
Sexually assaulted	2.7* (2.2–3.3)	1.5* (1.2–2.0)	1.5* (1.1–1.9)
Beaten up by spouse/romantic partner	2.4* (1.9–3.1)	1.2 (0.9–1.6)	1.3 (1.0–1.6)
Stalked	2.4* (1.9–3.1)	1.2 (0.9–1.6)	1.2 (0.9–1.6)
Traumatic event to loved one	2.6* (2.1–3.3)	1.3* (1.0–1.7)	1.4* (1.1–1.8)
Private event	2.5* (1.9–3.3)	1.5* (1.1–2.0)	1.5* (1.1–2.0)
Some other event	1.9* (1.4–2.7)	1.3 (0.9–1.8)	1.3 (0.9–1.8)
Accident/injuries			
Child with serious illness	2.2* (1.6–3.1)	1.6* (1.1–2.2)	1.5* (1.1–2.1)
Natural disaster	1.6* (1.2–2.1)	1.1 (0.8–1.5)	1.1 (0.8–1.4)
Life-threatening illness	1.7* (1.3–2.0)	1.2 (1.0–1.5)	1.2 (0.9–1.4)
Toxic chemical exposure	2.4* (1.7–3.3)	1.5* (1.1–2.2)	1.5* (1.1–2.1)
Other life threatening accident	2.0* (1.5–2.6)	1.2 (0.9–1.6)	1.2 (0.9–1.6)
Automobile accident	2.1* (1.7–2.5)	1.4* (1.1–1.7)	1.3* (1.1–1.6)
Other traumas			
Unexpected death of a loved one	1.6* (1.4–1.9)	1.1 (1.0–1.3)	1.1 (0.9–1.3)
Mugged or threatened with a weapon	2.0* (1.6–2.5)	1.3 (1.0–1.6)	1.2 (0.9–1.6)
Man-made disaster	1.8* (1.3–2.5)	1.0 (0.7–1.4)	1.0 (0.7–1.4)
Any traumatic events	3.1* (2.7–3.7)	NA	NA
Joint significance of all 29 traumatic events, $\chi^2$ (d.f.) <i>P</i>	NA	745.7* (29) <i>P</i> <0.001	128.4* (29) <i>P</i> <0.001
Differences in the ORs of the 29 traumatic events, $\chi^2$ (d.f.) <i>P</i>	NA	71.7* (29) <i>P</i> <0.001	64.6* (28) <i>P</i> <0.001
NA, not applicable			
a. Each lifetime traumatic event type was used as a predictor of psychotic experiences onset in separate discrete-time survival model controlling for age-cohorts, gender, person-year dummies, and country			
b. Model was estimated with dummy variables for all temporally prior traumatic events entered simultaneously including the controls specified above			
c. Model was estimated with dummy variables for type and number of traumatic events (starting at 2 traumatic events, 3 traumatic events, . . . , 5 or more traumatic events) entered simultaneously as predictors of psychotic experiences onset including the controls specified in (a)			
* <i>P</i> <0.05, 2-sided test			

<b>Table 2</b> Associations between number of temporally prior traumatic experiences and the subsequent onset of psychotic experiences		
	M2 – Multivariate number model <sup>a</sup> OR (95% CI)	M3 – Multivariate simple interactive model <sup>b</sup> OR (95% CI)
1 traumatic event	2.3* (1.9–2.8)	–
2 traumatic events	3.5* (2.8–4.4)	1.7* (1.4–2.1)
3 traumatic events	4.0* (3.1–5.1)	1.5* (1.1–2.0)
4 traumatic events	5.0* (3.8–6.7)	1.5* (1.1–2.2)
5 or more traumatic events	7.6* (6.1–9.5)	1.3 (0.8–2.1)
Joint significance of number-of-traumatic event measures $\chi^2$ (d.f.) <i>P</i>	355.9* (5) <i>P</i> <0.001	34.0* (4) <i>P</i> <0.001
<sup>a</sup> Model was estimated with dummy variables for all number of traumatic events without any information about type of traumatic events entered simultaneously controlling for age-cohorts, gender, person-year dummies, and country		
<sup>b</sup> Model was estimated with dummy variables for type and number of traumatic events (starting at 2 traumatic events, 3 traumatic events, . . . , 5 or more traumatic events) entered simultaneously as predictors of psychotic experiences onset including the controls specified in (a).		
* <i>P</i> <0.05, 2-sided test.		

**Table 3** Multivariate associations of traumatic experiences with the subsequent first onset of psychotic experiences, with and without adjustment for prior and intervening mental disorders

Type of traumatic experiences <sup>a</sup>	Model 4 – multivariate complex interactive model <sup>b</sup>	Model 5 (adjusted for PTSD) <sup>c</sup>	Model 6 (adjusted for mental disorders) <sup>d</sup>
Collective violence, marginal effects OR (95% CI)			
Civilian in war zone	0.4 (0.1–1.0)	0.4 (0.1–1.0)	0.4* (0.1–1.0)
Refugee	3.0 (0.8–11.0)	2.9 (0.8–10.7)	2.9 (0.8–10.3)
Civilian in region of terror	2.5 (0.6–9.8)	2.5 (0.6–9.8)	2.4 (0.6–9.6)
Kidnapped	1.3 (0.3–5.7)	1.3 (0.3–5.7)	1.2 (0.2–5.7)
Caused/witnessed bodily harm, marginal effects OR (95% CI)			
Combat experience	1.2 (0.2–5.7)	1.1 (0.2–5.7)	1.2 (0.2–5.9)
Accidentally caused serious injury or death	2.3 (0.7–7.7)	2.3 (0.7–7.7)	1.7 (0.6–4.5)
Saw atrocities	1.5 (0.5–4.7)	1.5 (0.5–4.7)	1.5 (0.5–4.6)
Witnessed death/dead body or saw someone seriously hurt	1.9* (1.4–2.5)	1.9* (1.4–2.5)	1.8* (1.3–2.5)
Interpersonal violence, marginal effects OR (95% CI)			
Beaten up by caregiver	2.0* (1.4–2.9)	2.0* (1.4–2.8)	1.9* (1.3–2.7)
Witnessed physical fight at home	1.8* (1.4–2.3)	1.8* (1.4–2.3)	1.8* (1.4–2.3)
Beaten up by someone else	1.9 (0.9–4.2)	1.9 (0.9–4.2)	1.8 (0.8–3.9)
Intimate partner/sexual violence, marginal effects OR (95% CI)			
Raped	4.9* (2.5–9.5)	4.8* (2.5–9.5)	4.3* (2.2–8.4)
Sexually assaulted	2.2* (1.4–3.6)	2.2* (1.3–3.5)	2.0* (1.3–3.3)
Beaten up by spouse/romantic partner	1.4 (0.8–2.6)	1.4 (0.8–2.6)	1.3 (0.7–2.3)
Stalked	3.4* (2.0–5.6)	3.4* (2.0–5.6)	3.0* (1.8–5.0)
Traumatic event to loved one	2.0* (1.1–3.5)	2.0* (1.1–3.5)	1.8* (1.0–3.3)
Private event	2.0 (0.9–4.5)	2.0 (0.9–4.5)	1.9 (0.9–4.2)
Some other event	1.5 (0.7–3.1)	1.5 (0.7–3.1)	1.4 (0.7–2.9)
Accident/injuries, marginal effects OR (95% CI)			
Child with serious illness	1.8 (0.7–4.5)	1.8 (0.7–4.5)	1.9 (0.8–4.6)
Natural disaster	1.2 (0.7–2.0)	1.2 (0.7–2.0)	1.2 (0.7–1.9)
Life-threatening illness	1.5* (1.0–2.3)	1.5* (1.0–2.3)	1.5 (1.0–2.3)
Toxic chemical exposure	1.5 (0.7–3.4)	1.5 (0.7–3.4)	1.5 (0.7–3.2)
Other life-threatening accident	2.1* (1.3–3.4)	2.1* (1.3–3.4)	2.0* (1.3–3.3)
Automobile accident	2.0* (1.4–3.0)	2.0* (1.4–3.0)	2.0* (1.3–2.9)
Other traumas, marginal effects OR (95% CI)			
Unexpected death of a loved one	1.5* (1.1–2.0)	1.5* (1.1–2.0)	1.5* (1.1–2.0)
Mugged or threatened with a weapon	1.9* (1.1–3.5)	1.9* (1.1–3.5)	1.8 (1.0–3.3)
Man-made disaster	1.2 (0.5–2.8)	1.2 (0.5–2.8)	1.2 (0.5–2.9)
Joint significance of all traumatic events, $\chi^2$ (d.f.) <i>P</i>	140.9* (29) <0.001	139.9* (29) <0.001	118.2* (29) <0.001
Number of other co-occurring traumatic events			
Marginal effects OR (95% CI)	1.1* (1.0–1.3)	1.1* (1.0–1.3)	1.1* (1.0–1.3)
$\chi^2$ (d.f.) <i>P</i>	5.1*(1) 0.024	5.2* (1) 0.023	5.2* (1) 0.023

a. The traumatic events: 'Relief worker in war zone' and 'Purposely injured, tortured, or killed someone' are not included in the table as the results are not applicable because of unstable estimates. Total person-years used for this multivariate model is 1 041 109.

b. Complex interactive model was estimated with dummy variables for type and number of temporally prior traumatic events and interaction terms for each type of traumatic events and number of other traumatic events including the controls: age-cohorts, gender, person-year dummies, and country.

c. Model specification in (b) and additionally control for temporally prior post-traumatic stress disorder (PTSD) only.

d. Model specification in (b) and additionally control for 21 temporally prior and intervening mental disorders (PTSD inclusive).

\**P* < 0.05, 2-sided test.

consistent with recent literature that traumatic events are associated with psychotic experiences in a dose-dependent fashion.<sup>3–6,8</sup> In addition, the study contributed two important findings: first, traumatic events were not equipotent with respect to the subsequent first onset of psychotic experiences. Second, comorbid mental disorders (including PTSD) did not account for the associations of traumatic events with psychotic experiences.

### Variation in the strength of association

With respect to the variation in the strength of association across traumatic event types, prior research has often failed to use statistical models that adequately address the intercorrelated nature of traumatic events. We constructed multivariable models that included a comprehensive list of 29 traumatic event types, and incorporated number of traumatic events per person, as well as interactive models. Similar to our earlier analysis of childhood adversities and psychotic experience onsets,<sup>2</sup> we found that not all

traumatic events had equivalent associations with subsequent first onset of psychotic experiences. For example, in the marginal effect model (model 4), we found that only 12 of the 29 traumatic event types were associated with an increased risk of subsequent psychotic experiences. Six of the ten traumatic events related to interpersonal violence, intimate partner, and sexual violence were associated with psychotic experiences (effect sizes ranged from 1.8 to 4.9 for 'witnessed physical fight at home' and 'raped', respectively). Those who endorsed 'raped' had fivefold increased odds of subsequent onset of psychotic experiences (OR = 4.9, 95% CI 2.5–9.5). The general pattern of findings indicates a complex interactive association between traumatic event types and number of traumatic events. In the complex interactive model (model 4), we also found that those who were exposed to more traumatic events were more likely to develop a first-onset psychotic experience regardless of type of traumatic events. This is consistent with the hypothesis that traumatic event exposure is a risk factor for psychosis,<sup>20</sup> and also lends additional support for the

important role of stress in aetiological pathways underpinning risk of psychotic experiences. Interestingly, in some models, ‘civilian or relief worker in war zone’ showed protective effects with respect to subsequent psychotic experiences onset. This is consistent with previous WMH survey studies with traumatic events, which suggest that respondents who had war experience may have a reduced risk of mental or physical disorders compared with the general population.<sup>21</sup> This may partly be because of a selection effect and/or to prior exposure promoting resilience.<sup>21</sup> In summary, the nature of the relationship between exposure to traumatic events and subsequent psychotic experiences is more nuanced than previously thought.

### Role of comorbid disorders

With respect to the explanatory effects of comorbid disorders, we found that the associations of traumatic events with psychotic experiences were mostly unchanged after adjustment for mental disorders that began prior to the onset of psychotic experiences. This adjustment included both confounding effects (i.e. mental disorders that both predicted subsequent occurrence of traumatic events and later onset of psychotic experiences) and mediating effects (i.e. traumatic events predicting subsequent onset of mental disorders that, in turn, are predictors of later psychotic experiences). No attempt was made to distinguish between the two types of effects. The finding that comorbid mental disorders do not account for the associations of traumatic events with subsequent psychotic experiences is consistent with our earlier findings that there was little or no influence of mental disorders on the associations of childhood adversities with psychotic experiences.<sup>2</sup> In particular, we found that PTSD did not appear to play a substantial role in accounting for the associations of traumatic events with psychotic experiences. This was in contrast to an earlier finding, where those with both exposure to traumatic events and associated PTSD were at greater risk of lifetime psychotic experiences.<sup>8</sup> However, our finding is consistent with the hypothesis that some (but not all) traumatic events may be ‘sufficient’ causes of subsequent psychotic experience onset (i.e. an intervening mental disorder is not required to trigger psychotic experiences). Of course, genetic factors and other unmeasured confounders could influence this association.

Even though the exact mechanisms linking traumatic events to psychotic experiences remain poorly understood, there is considerable recent interest in how socially mediated factors and traumatic events may influence the aetiology of psychotic experiences. Traumatic events may create a constitutional disposition for psychotic experiences that may include heightened vulnerability to stress,<sup>22</sup> elevated emotional reactivity and dysregulation<sup>21,23</sup> and deficits in cognitive control.<sup>24</sup> For example, it is feasible that psychopathology emerges as the end product of distress and anxiety because of maladaptive stress and emotional reactions of traumatic events, which in turn may lead to dopamine sensitisation. These pathways may influence psychosis vulnerability.<sup>25</sup> Stress-related changes in the hypothalamic axis could also have an impact on brain circuitry via epigenetic mechanisms, leading to vulnerability to subsequent psychotic experiences.<sup>26</sup>

### Limitations

Although the current study has many strengths (for example range of traumatic event types, large sample size, range of countries, uniform methodology for data collection), it also has several limitations that deserve comment. We excluded those who screened positive for possible psychotic disorders based on self-reporting having received a psychosis diagnosis or having used

antipsychotic medications to treat the reported psychotic experiences. However, it is possible that some respondents who reported psychotic experiences had an untreated psychotic disorder. We also relied on retrospective reports about age at onset, which might have led to a recall bias. However, we note that several prospective studies have confirmed the association between traumatic events and subsequent psychotic experiences.<sup>4,5</sup> Although we had age at exposure for each type of traumatic event, we could not define the pattern of exposure across the life course (for example we could not identify those who only had childhood exposure *v.* those that had both childhood and adult exposure). We failed to control for all plausible correlates of both traumatic event exposure and psychotic experiences, as we wanted to focus on comorbid mental disorders and we were concerned that other plausible correlates (such as socioeconomic status and education) could be either causes or consequences of both traumatic event exposure and psychotic experiences. Finally, while our findings are consistent with the hypothesis that particular types of traumatic event exposure may contribute to the development of subsequent psychotic experiences, the surveys are cross-sectional and thus unable to determine causal pathways.

### Implications

Despite these limitations, our study found that many of the traumatic events (11 out of 29) were independently associated with subsequent psychotic experiences in a large, community-based transnational study that were not mediated by mental disorders. The robust literature now linking traumatic events and subsequent psychotic experiences may enrich the epidemiological landscape of psychotic experiences, and can provide important clues to the aetiology and pathogenesis of psychotic experiences.

### Funding

The World Health Organization World Mental Health (WMH) Survey Initiative is supported by the National Institute of Mental Health (NIMH; R01 MH070884), the John D. and Catherine T. MacArthur Foundation, the Pfizer Foundation, the US Public Health Service (R13-MH066849, R01-MH069864, and R01 DA016558), the Fogarty International Center (FIRCA R03-TW006481), the Pan American Health Organization, Eli Lilly and Company, Ortho-McNeil Pharmaceutical, GlaxoSmithKline, and Bristol-Myers Squibb. None of the funders had any role in the design, analysis, interpretation of results, or preparation of this paper. The views and opinions expressed in this report are those of the authors and should not be construed to represent the views of the World Health Organization, other sponsoring organisations, agencies, or governments.

The Colombian National Study of Mental Health (NSMH) is supported by the Ministry of Social Protection. The ESEMeD project is funded by the European Commission (Contracts QLGS-1999-01042; SANCO 2004123, and EAHC 20081308), the Piedmont Region (Italy), Fondo de Investigación Sanitaria, Instituto de Salud Carlos III, Spain (FIS 00/0028), Ministerio de Ciencia y Tecnología, Spain (SAF 2000-158-CE), Departament de Salut, Generalitat de Catalunya, Spain, Instituto de Salud Carlos III (CIBER CB06/02/0046, RETICS RD06/0011 REM-TAP), and other local agencies and by an unrestricted educational grant from GlaxoSmithKline. The Lebanese Evaluation of the Burden of Ailments and Needs of the Nation (L.E.B.A.N.O.N.) is supported by the Lebanese Ministry of Public Health, the WHO (Lebanon), National Institute of Health / Fogarty International Center (R03 TW006481-01), anonymous private donations to IDRAAC, Lebanon, and unrestricted grants from, Algorithm, AstraZeneca, Benta, Bella Pharma, Eli Lilly, Glaxo Smith Kline, Lundbeck, Novartis, Servier, Phénicia, UPO. The Mexican National Comorbidity Survey (MNCS) is supported by The National Institute of Psychiatry Ramon de la Fuente (INPRFMDIES 4280) and by the National Council on Science and Technology (CONACyT-G30544- H), with supplemental support from the PanAmerican Health Organization (PAHO). Te Rau Hinengaro: The New Zealand Mental Health Survey (NZMHS) is supported by the New Zealand Ministry of Health, Alcohol Advisory Council, and the Health Research Council. The Nigerian Survey of Mental Health and Wellbeing (NSMHW) is supported by the WHO (Geneva), the WHO (Nigeria), and the Federal Ministry of Health, Abuja, Nigeria. The Peruvian World Mental Health Study was funded by the National Institute of Health of the Ministry of Health of Peru. The Portuguese Mental Health Study was carried out by the Department of Mental Health, Faculty of Medical Sciences, NOVA University of Lisbon, with collaboration of the Portuguese Catholic University, and was funded by Champalimaud Foundation, Gulbenkian Foundation, Foundation for Science and Technology (FCT) and Ministry of Health. The Romania WMH study projects ‘Policies in Mental Health Area’ and ‘National Study regarding Mental Health and Services Use’ were carried out by National School of Public Health & Health Services Management (former National Institute for Research & Development in Health, present National School of Public Health Management & Professional Development, Bucharest), with technical support of Metro Media Transilvania, the National Institute of Statistics – National Centre for Training in Statistics, SC. Cheyenne Services SRL, Statistics Netherlands and were funded by Ministry of Public Health (former Ministry of Health) with supplemental support of Eli Lilly Romania SRL. The São Paulo Megacity Mental Health Survey is supported by the State of São Paulo



Research Foundation (FAPESP) Thematic Project Grant 03/00204-3The US National Comorbidity Survey Replication (NCS-R) is supported by the National Institute of Mental Health (NIMH; U01-MH60220) with supplemental support from the National Institute of Drug Abuse (NIDA), the Substance Abuse and Mental Health Services Administration (SAMHSA), the Robert Wood Johnson Foundation (RWJF; Grant 044708), and the John W. Alden Trust. J.J.M. received the John Cade Fellowship APP1056929 from the National Health and Medical Research Council, and Niels Bohr Professorship from the Danish National Research Foundation.

## Acknowledgements

A complete list of all within-country and cross-national WMH publications can be found at <http://www.hcp.med.harvard.edu/wmh/>. We thank the staff of the WMH Data Collection and Data Analysis Coordination Centres for assistance with instrumentation, fieldwork, and consultation on data analysis.

**John J. McGrath**, MD, PhD, Queensland Centre for Mental Health Research, and Queensland Brain Institute, University of Queensland, and National Centre for Register-based Research, Aarhus BSS, Aarhus University, Aarhus, Denmark; **Sukanta Saha**, PhD, Queensland Centre for Mental Health Research, and Queensland Brain Institute, University of Queensland, Australia; **Carmen C. W. Lim**, MSc, Queensland Brain Institute, The University of Queensland, St. Lucia, Queensland, Australia; **Sergio Aguilar-Gaxiola**, MD, PhD, Center for Reducing Health Disparities, UC Davis Health System, Sacramento, California, USA; **Jordi Alonso**, MD, PhD, Health Services Research Unit, IMIM-Hospital del Mar Medical Research Institute, Barcelona, Spain, Pompeu Fabra University (UPF), Barcelona, Spain and CIBER en Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain; **Laura H. Andrade**, MD, PhD, Section of Psychiatric Epidemiology – LIM 23, Institute of Psychiatry, University of São Paulo Medical School, São Paulo, Brazil; **Evelyn J. Bromet**, PhD, Department of Psychiatry, Stony Brook University School of Medicine, Stony Brook, New York, USA; **Ronny Bruffaerts**, PhD, Universitair Psychiatrisch Centrum – Katholieke Universiteit Leuven (UPC-KUL), Campus Gasthuisberg, Leuven, Belgium; **José M. Caldas de Almeida**, MD, PhD, Chronic Diseases Research Center (CEDOC) and Department of Mental Health, Faculdade de Ciências Médicas, Universidade Nova de Lisboa, (Campo dos Mártires da Pátria), Lisbon, Portugal; **Graca Cardoso**, MD, PhD, Lisbon Institute of Global Mental Health and Chronic Diseases Research Center (CEDOC), Nova Medical School, Universidade Nova de Lisboa, Lisboa, Portugal; **Giovanni de Girolamo**, MD, Unit of Epidemiological and Evaluation Psychiatry, Istituti di Ricovero e Cura a Carattere Scientifico (IRCCS) – St. John of God Clinical Research Centre, Brescia, Italy; **John Fayyad**, MD, Institute for Development, Research, Advocacy & Applied Care (IDRAAC), Beirut, Lebanon; **Silvia Florescu**, MD, PhD, National School of Public Health, Management and Professional Development, Bucharest, Romania; **Oye Gureje**, MD, DSc, FRCPsych, Department of Psychiatry, University College Hospital, Ibadan, Nigeria; **Josep M. Haro**, MD, PhD, Parc Sanitari Sant Joan de Déu, CIBERSAM, Universitat de Barcelona, Barcelona, Spain; **Norito Kawakami**, MD, DMSc, Department of Mental Health, School of Public Health, The University of Tokyo, Tokyo, Japan; **Karestan C. Koenen**, PhD, Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA; **Viviane Kovess-Masfety**, MD, PhD, Ecole des Hautes Etudes en Santé Publique (EHESP), EA 4057 Paris Descartes University, Paris, France; **Sing Lee**, MBBS, Department of Psychiatry, Chinese University of Hong Kong, Tai Po, Hong Kong; **Jean-Pierre Lepine**, MD, Hôpital Lariboisière- Fernand Widal, Assistance Publique Hôpitaux de Paris, Universités Paris Descartes-Paris Diderot and INSERM UMR-S 1144, Paris, France; **Katie A. McLaughlin**, PhD, Department of Psychology, University of Washington, Seattle, Washington, USA; **Maria E. Medina-Mora**, PhD, National Institute of Psychiatry Ramón de la Fuente, Mexico City, Mexico; **Fernando Navarro-Mateu**, MD, PhD, IMIB-Arixaca, CIBERESP-Murcia, Subdirección General de Salud Mental y Asistencia Psiquiátrica, Servicio Murciano delusional experiences Salud, El Palmar (Murcia), Spain; **Akin Ojagbemi**, PhD, College of Medicine, University of Ibadan; University College Hospital, Ibadan, Nigeria; **Jose Posada-Villa**, MD, Colegio Mayor de Cundinamarca University, Faculty of Social Sciences, Bogota, Colombia; **Nancy Sampson**, BA, Department of Health Care Policy, Harvard Medical School, Boston, Massachusetts, USA; **Kate M. Scott**, PhD, Department of Psychological Medicine, University of Otago, Dunedin, Otago, New Zealand; **Hisateru Tachimori**, PhD, National Institute of Mental Health, National Center for Neurology and Psychiatry, Tokyo, Japan; **Margreet ten Have**, PhD, Trimbos-instituut, Netherlands Institute of Mental Health and Addiction, Netherlands; **Kenneth S. Kendler**, MD, Department of Psychiatry, Virginia Commonwealth University, USA; **Ronald C. Kessler**, PhD, Department of Health Care Policy, Harvard Medical School, Boston, Massachusetts, USA

**Correspondence:** John McGrath, Queensland Brain Institute, The University of Queensland, St Lucia, Queensland 4076, Australia. Email: [j.mcgrath@uq.edu.au](mailto:j.mcgrath@uq.edu.au)

First received 4 May 2017, final revision 21 Jun 2017, accepted 26 Jul 2017

## Appendix

The WHO World Mental Health Survey collaborators are: Sergio Aguilar-Gaxiola, MD, PhD, Ali Al-Hamzawi, MD, Mohammed Salih Al-Kaisy, MD, Jordi Alonso, MD, PhD, Laura Helena Andrade, MD, PhD, Corina Benjet, PhD, Guilherme Borges, ScD, Evelyn J. Bromet, PhD, Ronny Bruffaerts, PhD, Brendan Bunting, PhD, Jose Miguel Caldas de Almeida, MD, PhD, Graca Cardoso, MD, PhD, Somnath Chatterji, MD, Alfredo H. Cia, MD, Louisa Degenhardt, PhD, Koen Demyttenaere, MD, PhD, John Fayyad, MD, Silvia Florescu, MD, PhD, Giovanni de Girolamo, MD, Oye Gureje,

MD, DSc, FRCPsych, Josep Maria Haro, MD, PhD, Yanling He, MD, Hristo Hinkov, MD, PhD, Chi-yi Hu, MD, PhD, Yueqin Huang, MD, MPH, PhD, Peter de Jonge, PhD, Aimee Nasser Karam, PhD, Elie G. Karam, MD, Norito Kawakami, MD, DMSc, Ronald C. Kessler, PhD, Andrzej Kiejna, MD, PhD, Viviane Kovess-Masfety, MD, PhD, Sing Lee, MB, BS, Jean-Pierre Lepine, MD, Daphna Levinson, PhD, John McGrath, MD, PhD, Maria Elena Medina-Mora, PhD, Jacek Moskalewicz, PhD, Fernando Navarro-Mateu, MD, PhD, Beth-Ellen Pennell, MA, Marina Piazza, MPH, ScD, Jose Posada-Villa, MD, Kate M. Scott, PhD, Tim Slade, PhD, Juan Carlos Stagnaro, MD, PhD, Dan J. Stein, FRCPC, PhD, Margreet ten Have, PhD, Yolanda Torres, MPH, Dra.HC, Maria Carmen Viana, MD, PhD, Harvey Whiteford, MBBS, PhD, David R. Williams, MPH, PhD, Bogdan Wojtyniak, ScD.

## References

- Trotta A, Murray RM, Fisher HL. The impact of childhood adversity on the persistence of psychotic symptoms: a systematic review and meta-analysis. *Psychol Med* 2015; **45**: 2481–98.
- McGrath JJ, McLaughlin KA, Saha S, Aguilar-Gaxiola S, Al-Hamzawi A, Alonso J, et al. The association between childhood adversities and subsequent first onset of psychotic experiences: a cross-national analysis of 23 998 respondents from 17 countries. *Psychol Med* 2017; **47**: 1230–45.
- Beards S, Gayer-Anderson C, Borges S, Dewey ME, Fisher HL, Morgan C. Life events and psychosis: a review and meta-analysis. *Schizophr Bull* 2013; **39**: 740–7.
- Kelleher I, Harley M, Lynch F, Arseneault L, Fitzpatrick C, Cannon M. Associations between childhood trauma, bullying and psychotic symptoms among a school-based adolescent sample. *Br J Psychiatry* 2008; **193**: 378–82.
- Spauwen J, Krabbendam L, Lieb R, Wittchen H-U, van Os J. Impact of psychological trauma on the development of psychotic symptoms: relationship with psychosis proneness. *Br J Psychiatry* 2006; **188**: 527–33.
- Saha S, Varghese D, Slade T, Degenhardt L, Mills K, McGrath J, et al. The association between trauma and delusional-like experiences. *Psychiatry Res* 2011; **189**: 259–64.
- Jenkins R, Mbatia J, Singleton N, White B. Prevalence of psychotic symptoms and their risk factors in urban Tanzania. *Int J Environ Res Public Health* 2010; **7**: 2514–25.
- Scott J, Chant D, Andrews G, Martin G, McGrath J. Association between trauma exposure and delusional experiences in a large community-based sample. *Br J Psychiatry* 2007; **190**: 339–43.
- Kessler RC, McLaughlin KA, Koenen KC, Petukhova M, Hill ED. The importance of secondary trauma exposure for post-disaster mental disorder. *Epidemiol Psychiatr Sci* 2012; **21**: 35–45.
- Green JG, McLaughlin KA, Berglund PA, Gruber MJ, Sampson NA, Zaslavsky AM, et al. Childhood adversities and adult psychiatric disorders in the national comorbidity survey replication I: associations with first onset of DSM-IV disorders. *Arch Gen Psychiatry* 2010; **67**: 113–23.
- Karam EG, Friedman MJ, Hill ED, Kessler RC, McLaughlin KA, Petukhova M, et al. Cumulative traumas and risk thresholds: 12-month PTSD in the World Mental Health (WMH) surveys. *Depress Anxiety* 2014; **31**: 130–42.
- McGrath JJ, Saha S, Al-Hamzawi A, Andrade L, Benjet C, Bromet EJ, et al. The bidirectional associations between psychotic experiences and DSM-IV mental disorders. *Am J Psychiatry* 2016; **173**: 997–1006.
- Soosay I, Silove D, Bateman-Steel C, Steel Z, Bebbington P, Jones PB, et al. Trauma exposure, PTSD and psychotic-like symptoms in post-conflict Timor Leste: an epidemiological survey. *BMC Psychiatry* 2012; **12**: 229.
- Kessler RC, Ustun TB. The World Mental Health (WMH) Survey Initiative Version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). *Int J Methods Psychiatr Res* 2004; **13**: 93–121.
- Kessler RC, Ustun TB. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. Cambridge University Press, 2008.
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorder (4th edn) (DSM-IV)*. APA, 1994.
- World Health Organization. *The ICD-10 Classification of Mental and Behavioural Disorders: Clinical Descriptions and Diagnostic Guidelines*. WHO, 1992.
- Haro JM, Arbabzadeh-Bouchez S, Brugh TS, de Girolamo G, Guyer ME, Jin R, et al. Concordance of the Composite International Diagnostic Interview Version 3.0 (CIDI 3.0) with standardized clinical assessments in the WHO World Mental Health surveys. *Int J Methods Psychiatr Res* 2006; **15**: 167–80.

- 19 Benjet C, Bromet E, Karam EG, Kessler RC, McLaughlin KA, Ruscio AM, et al. The epidemiology of traumatic event exposure worldwide: results from the World Mental Health Survey Consortium. *Psychol Med* 2016; **46**: 327–43.
- 20 Freedman R. Investigating trauma as a risk factor for psychosis. *Schizophr Bull* 2017; **43**: 1–2.
- 21 Liu H, Petukhova MV, Sampson NA, Aguilar-Gaxiola S, Alonso J, Andrade LH, et al. Association of DSM-IV posttraumatic stress disorder with traumatic experience type and history in the World Health Organization World Mental Health Surveys. *JAMA Psychiatry* 2017; **74**: 270–81.
- 22 McLaughlin KA, Kubzansky LD, Dunn EC, Waldinger R, Vaillant G, Koenen KC. Childhood social environment, emotional reactivity to stress, and mood and anxiety disorders across the life course. *Depress Anxiety* 2010; **27**: 1087–94.
- 23 McCrory EJ, De Brito SA, Sebastian CL, Mechelli A, Bird G, Kelly PA, et al. Heightened neural reactivity to threat in child victims of family violence. *Curr Biol* 2011; **21**: R947–8.
- 24 DePrince AP, Weinzierl KM, Combs MD. Executive function performance and trauma exposure in a community sample of children. *Child Abuse Negl* 2009; **33**: 353–61.
- 25 Kapur S. Psychosis as a state of aberrant salience: a framework linking biology, phenomenology, and pharmacology in schizophrenia. *Am J Psychiatry* 2003; **160**: 13–23.
- 26 Barker V, Gumley A, Schwannauer M, Lawrie SM. An integrated biopsychosocial model of childhood maltreatment and psychosis. *Br J Psychiatry* 2015; **206**: 177–80.





Data supplement to McGrath et al. Trauma and psychotic experiences: transnational data from the World Mental Health Survey. Br J Psychiatry doi: 10.1192/bjp.bp.117.205955

**Table DS1 World Mental Health (WMH) sample characteristics by World Bank income categories, and sample for psychotic experiences (PEs)**

Country by income category	Survey <sup>a</sup>	Sample characteristics	Field dates	Age range	Sample size		Response rate <sup>c</sup>
					Part I	PEs sample	
<b>Low and lower middle income countries</b>							
Colombia	NSMH	All urban areas of the country	2003	18-65	4426	722	87.7
Nigeria	NSMHW	21 of the 36 states in the country	2002-3	18-100	6752	1417	79.3
Peru	EMSMP	5 urban areas (approximately 38% of the total national population)	2004-5	18-65	3930	530	90.2
<b>Upper-middle income countries</b>							
Brazil - São Paulo	São Paulo Megacity	São Paulo metropolitan area	2005-7	18-93	5037	2922	81.3
Lebanon	LEBANON	Nationally representative	2002-3	18-94	2857	1029	70
Mexico	M-NCS	All urban areas of the country	2001-2	18-65	5782	715	76.6
Romania	RMHS	Nationally representative	2005-6	18-96	2357	2357	70.9
<b>High-income countries</b>							
Belgium	ESEMeD	Nationally representative	2001-2	18-95	2419	319	50.6
France	ESEMeD	Nationally representative	2001-2	18-97	2894	301	45.9
Germany	ESEMeD	Nationally representative	2002-3	18-95	3555	408	57.8
Italy	ESEMeD	Nationally representative	2001-2	18-100	4712	617	71.3
New Zealand <sup>d</sup>	NZMHS	Nationally representative	2003-4	18-98	12790	7263	73.3
Portugal	NMHS	Nationally representative	2008-9	18-81	3849	2053	57.3
Spain	ESEMeD	Nationally representative	2001-2	18-98	5473	1159	78.6
The Netherlands	ESEMeD	Nationally representative	2002-3	18-95	2372	348	56.4
The United States	NCS-R	Nationally representative	2002-3	18-99	9282	2304	70.9
<b>All countries combined</b>					<b>78487</b>	<b>24464</b>	<b>70.5</b>

<sup>a</sup> NSMH (The Colombian National Study of Mental Health); NSMHW (The Nigerian Survey of Mental Health and Wellbeing); EMSMP (La Encuesta Mundial de Salud Mental en el Peru); LEBANON (Lebanese Evaluation of the Burden of Ailments and Needs of the Nation); M-NCS (The Mexico National Comorbidity Survey); RMHS (Romania Mental Health Survey); ESEMeD (The European Study Of The Epidemiology Of Mental Disorders); NZMHS (New Zealand Mental Health Survey); NMHS (Portugal National Mental Health Survey); NCS-R (The US National Comorbidity Survey Replication).

<sup>b</sup> Brazil, New Zealand and Romania did not have an age restricted sample. All other countries, with the exception of Nigeria (which was age restricted to ≤ 39) were age restricted to ≤ 44.

<sup>c</sup> The response rate is calculated as the ratio of the number of households in which an interview was completed to the number of households originally sampled, excluding from the denominator households known not to be eligible either because of being vacant at the time of initial contact or because the residents were unable to speak the designated languages of the survey. The weighted average response rate is 70.5%.

<sup>d</sup> For the purposes of cross-national comparisons we limit the sample to those 18+.

**Table DS2 Six CIDI Psychotic experiences types in six European (ESEMed<sup>a</sup>) sites (Belgium, France, Germany, Italy, Netherlands, Spain)**

<b>Item</b>	<b>Type</b>	<b>Description</b>
<b>A. Saw a vision (Visual hallucinations)</b>	<b>1</b>	Did you ever see something that wasn't really there that other people could not see? Please do not include any times when you were dreaming or half-asleep or under the influence of alcohol or drugs.
<b>B. Heard voices (Auditory hallucinations)</b>	<b>2</b>	Did you ever hear things that other people said did not exist, like strange voices coming from inside your head talking to you or about you, or voices coming out of the air when there was no one around. Please do not include any times when you were dreaming or half-asleep or under the influence of alcohol or drugs.
<b>C. Thought insertion/withdrawal</b>	<b>3</b>	Did you ever believe that some mysterious force was inserting many different strange thoughts -- that were definitely not your own thoughts -- directly into your head by means of x-rays or laser beams or other methods?
<b>D. Mind control/passivity</b>	<b>4</b>	Did you ever feel that your mind had been taken over by strange forces with laser beams or other methods that were making you do things you did not choose to do. Again, do not include times when you were dreaming or under the influence of alcohol or drugs.
<b>E. Ideas of reference</b>	<b>5</b>	Did you ever believe that some strange force was trying to communicate directly with you by sending special signs or signals that you could understand but that no one else could understand. Sometimes this happens by special signs coming through the radio or television.
<b>F. Plot to harm/follow</b>	<b>6</b>	Did you ever believe that there was an unjust plot going on to harm you or to have people follow you that your family and friends did not believe existed?

<sup>a</sup> ESEMed = European Study of the Epidemiology of Mental Disorders

**Table DS3 Six CIDI Psychotic experiences types in 10 non-ESEMed sites (Colombia, Lebanon, Mexico, Brazil, Nigeria, Peru, Portugal, Romania, USA and New Zealand)**

<b>Item</b>	<b>Type</b>	<b>Description</b>
<b>A. Saw a vision (Visual hallucinations)</b>	<b>1</b>	Did you ever see something that other people who were there could not see?
	<b>1a</b>	Did this every happen when you were not dreaming, not half-asleep, and not under the influence of alcohol or drugs?
<b>B. Hearing voices (Auditory hallucinations)</b>	<b>2</b>	Did you ever hear things that other people said did not exist, like strange voices coming from inside your head talking to you or about you, or voices coming out of the air when there was no one around.
	<b>2a</b>	Did this every happen when you were not dreaming, not half-asleep, and not under the influence of alcohol or drugs?
<b>C. Thought insertion/withdrawal</b>	<b>3</b>	Did you ever believe that some mysterious force was inserting many different strange thoughts -- that were definitely not your own thoughts -- directly into your head by means of x-rays or laser beams or other methods?
	<b>3a</b>	Did this every happen when you were not dreaming, not half-asleep, and not under the influence of alcohol or drugs?
<b>D. Mind control/passivity</b>	<b>4</b>	Did you ever feel that your mind had been taken over by strange forces with laser beams or other methods that were making you do things you did not choose to do.
	<b>4a</b>	Did this every happen when you were not dreaming, not half-asleep, and not under the influence of alcohol or drugs?
<b>E. Ideas of reference</b>	<b>5</b>	Did you ever believe that some strange force was trying to communicate directly with you by sending special signs or signals that you could understand but that no one else could understand. Sometimes this happens by special signs coming through the radio or television.
	<b>5a</b>	Did this every happen when you were not dreaming, not half-asleep, and not under the influence of alcohol or drugs?
<b>F. Plot to harm/follow</b>	<b>6</b>	Did you ever believe that there was an unjust plot going on to harm you or to have people follow you that your family and friends did not believe existed?
	<b>6a</b>	Did this every happen when you were not dreaming, not half-asleep, and not under the influence of alcohol or drugs?

Note: For the assessment of psychotic experiences we included items 1a, 2a, 3a, 4a, 5a, and 6a. Similarly for the assessment of hallucination types we included types 1a and 2a, and for the assessment of delusional experiences we included types 3a, 4a, 5a, and 6a.



**Table DS4 21 DSM-IV mental disorders across 16 WMH sites**

---

<b>A. Mood disorders</b>	Major depressive disorder Bipolar disorder (Bipolar I, II, Subthreshold)
<b>B. Anxiety disorders</b>	Panic disorder Generalized anxiety disorder Specific phobia Social phobia Agoraphobia without panic Post-traumatic stress disorder Separation anxiety disorder (Child) Separation anxiety disorder (Adult)
<b>C. Behaviour disorders</b>	Intermittent explosive disorder Attention deficit disorder Oppositional defiant disorder Conduct disorder
<b>D. Eating disorders</b>	Anorexia nervosa Bulimia nervosa Binge eating disorder
<b>E. Substance-use disorders</b>	Alcohol abuse Alcohol dependence Drug abuse Drug dependence

---

Table D55 Prevalence of traumatic experiences (TE) among respondents with and without lifetime psychotic experiences (PE) (n = 24,464)

Type of traumatic experiences	Total Sample			Without PE			With PE			Respondents endorsing both lifetime TE and PE							
	n	% <sup>b</sup>	S.E	n	% <sup>b</sup>	S.E	n	% <sup>b</sup>	S.E	TE prior to PE onset		TE in the same year as PE onset		PE prior to TE onset		Goodness-of-fit test for equal proportion <sup>a</sup>	
										% <sup>b</sup>	S.E	% <sup>b</sup>	S.E	% <sup>b</sup>	S.E	$\chi^2$	[p]
<b>I. Collective violence</b>																	
Civilian in war zone <sup>c</sup>	1423	6.4	0.3	1331	6.5	0.3	92	4.6	0.8	70.9	7.3	2.2	1.4	26.9	7.2	111.6*	<.001
Refugee <sup>c</sup>	714	3.0	0.2	666	3.0	0.3	48	2.7	0.6	76.2	7.4	1.7	1.6	22.1	7.2	22.6*	<.001
Civilian in region of terror <sup>c</sup>	699	3.3	0.2	622	3.2	0.2	77	4.5	0.7	61.4	7.6	5.0	2.7	33.6	7.3	12.9*	<.001
Kidnapped	370	1.1	0.1	289	1.0	0.1	81	2.9	0.4	43.8	7.5	5.1	2.2	51.2	7.6	2.2	0.137
Relief worker in war zone <sup>c</sup>	242	1.2	0.1	225	1.2	0.1	17	0.9	0.3	40.3	14.5	9.9	9.3	49.8	15.4	- <sup>f</sup>	- <sup>f</sup>
<b>Any collective violence</b>	<b>2454</b>	<b>9.7</b>	<b>0.3</b>	<b>2214</b>	<b>9.7</b>	<b>0.3</b>	<b>240</b>	<b>9.9</b>	<b>0.9</b>	<b>62.6</b>	<b>4.4</b>	<b>3.4</b>	<b>1.2</b>	<b>34.0</b>	<b>4.4</b>	<b>24.5*</b>	<b>&lt;.001</b>
<b>II. Caused/witnessed bodily harm</b>																	
Purposely injured, tortured, or killed someone <sup>d</sup>	153	0.7	0.1	121	0.7	0.1	32	1.8	0.5	53.4	12.8	2.8	2.9	43.7	12.7	- <sup>f</sup>	- <sup>f</sup>
Combat experience <sup>c</sup>	578	2.9	0.2	512	2.8	0.2	66	5.1	0.9	52.3	8.7	1.4	1.1	46.3	8.7	230.7*	<.001
Accidentally caused serious injury or death	457	1.5	0.1	375	1.4	0.1	82	3.2	0.4	65.4	6.3	3.1	2.4	31.4	6.1	61.8*	<.001
Saw atrocities	992	4.0	0.2	853	3.8	0.2	139	6.4	0.8	46.3	6.1	7.8	3.7	46.0	7.1	0.0	0.972
Witnessed death/dead body or saw someone seriously hurt	6422	24.4	0.4	5475	23.1	0.4	947	41.8	1.5	52.5	2.7	4.9	0.9	42.6	2.7	5.9*	0.015
<b>Any caused/witnessed bodily harm</b>	<b>6981</b>	<b>26.8</b>	<b>0.4</b>	<b>5980</b>	<b>25.6</b>	<b>0.4</b>	<b>1001</b>	<b>43.7</b>	<b>1.5</b>	<b>54.3</b>	<b>2.6</b>	<b>4.8</b>	<b>0.8</b>	<b>40.8</b>	<b>2.5</b>	<b>11.2*</b>	<b>0.001</b>
<b>III. Interpersonal violence</b>																	
Beaten up by caregiver	2948	8.2	0.2	2417	7.4	0.2	531	19.4	1.1	87.8	1.9	4.2	1.1	8.0	1.7	223.3*	<.001
Witnessed physical fight at home <sup>e</sup>	3238	12.0	0.3	2704	11.2	0.3	534	22.4	1.3	84.0	2.7	4.8	1.4	11.3	2.5	233.2*	<.001
Beaten up by someone else	1650	5.4	0.2	1381	5.1	0.2	269	9.6	0.8	60.6	4.0	9.1	2.4	30.2	3.6	39.1*	<.001
<b>Any interpersonal violence</b>	<b>5733</b>	<b>17.9</b>	<b>0.3</b>	<b>4811</b>	<b>16.6</b>	<b>0.3</b>	<b>922</b>	<b>35.2</b>	<b>1.5</b>	<b>84.8</b>	<b>1.8</b>	<b>3.9</b>	<b>0.8</b>	<b>11.3</b>	<b>1.7</b>	<b>392.8*</b>	<b>&lt;.001</b>
<b>IV. Intimate partner/sexual violence</b>																	
Raped	1485	3.5	0.2	1140	2.9	0.1	345	12.1	1.0	63.8	4.0	6.7	1.6	29.5	3.8	72.1*	<.001
Sexually assaulted	2179	6.1	0.2	1727	5.4	0.2	452	15.4	1.0	68.9	3.3	6.7	1.7	24.4	3.1	169.7*	<.001
Beaten up by spouse/romantic partner	1796	4.3	0.2	1469	3.9	0.2	327	10.2	0.8	46.6	4.1	8.1	2.1	45.3	4.2	0.0	0.842
Stalked	1851	5.7	0.2	1435	5.0	0.2	416	14.8	1.0	38.7	3.6	8.6	3.0	52.8	3.7	9.8*	0.002
Traumatic event to loved one	1966	6.4	0.2	1572	5.8	0.2	394	14.9	1.0	38.7	3.6	5.8	1.5	55.5	3.5	14.4*	<.001
Private event	1451	4.4	0.2	1165	4.0	0.2	286	11.0	0.9	44.6	4.4	6.6	2.2	48.8	4.3	0.6	0.435
Some other event	1241	4.2	0.2	1006	3.8	0.2	235	9.3	0.9	32.6	4.5	3.0	1.1	64.4	4.7	47.3*	<.001
<b>Any intimate partner/sexual violence</b>	<b>7257</b>	<b>23.1</b>	<b>0.4</b>	<b>6028</b>	<b>21.3</b>	<b>0.4</b>	<b>1229</b>	<b>47.6</b>	<b>1.5</b>	<b>60.1</b>	<b>2.1</b>	<b>7.0</b>	<b>1.0</b>	<b>33.0</b>	<b>1.9</b>	<b>57.5*</b>	<b>&lt;.001</b>
<b>V. Accident/injuries</b>																	
Child with serious illness	2366	7.9	0.2	2004	7.4	0.2	362	15.7	1.1	30.7	3.8	7.2	1.8	62.0	4.1	29.8*	<.001
Natural disaster <sup>c</sup>	1946	8.3	0.3	1709	7.9	0.3	237	13.9	1.3	57.2	5.3	4.2	2.4	38.5	5.2	6.6*	0.010
Life-threatening illness	3627	12.7	0.3	3174	12.3	0.3	453	17.8	1.1	42.7	3.2	4.2	1.0	53.1	3.3	6.6*	0.010
Toxic chemical exposure	1070	4.5	0.2	884	4.1	0.2	186	9.4	0.9	43.9	5.4	5.6	2.6	50.5	5.4	1.0	0.310
Other life threatening accident	1774	6.7	0.2	1524	6.4	0.2	250	10.9	0.9	48.4	4.4	6.5	1.8	45.1	4.3	0.4	0.527
Automobile accident	4089	15.1	0.3	3503	14.4	0.3	586	25.5	1.3	47.7	3.0	5.6	1.5	46.7	3.1	0.0	0.846
<b>Any accident/injuries</b>	<b>10125</b>	<b>37.8</b>	<b>0.5</b>	<b>8873</b>	<b>36.6</b>	<b>0.5</b>	<b>1252</b>	<b>53.8</b>	<b>1.6</b>	<b>51.9</b>	<b>2.1</b>	<b>6.3</b>	<b>1.0</b>	<b>41.8</b>	<b>2.2</b>	<b>7.1*</b>	<b>0.008</b>
<b>VI. Other traumas</b>																	
Unexpected death of a loved one	8655	31.3	0.4	7566	30.2	0.5	1089	45.6	1.7	43.0	2.1	8.7	1.2	48.3	2.2	2.1	0.146

Mugged or threatened with a weapon	4134	<b>15.3</b>	0.4	3467	<b>14.3</b>	0.4	667	<b>29.9</b>	1.8	<b>40.2</b>	2.9	<b>6.1</b>	1.3	<b>53.6</b>	2.9	<b>11.9*</b>	0.001
Man-made disaster	1022	<b>3.7</b>	0.2	846	<b>3.5</b>	0.2	176	<b>7.2</b>	0.7	<b>44.4</b>	5.3	<b>3.6</b>	1.3	<b>52.0</b>	5.3	<b>1.3</b>	0.256
<b>Any other traumas</b>	<b>11061</b>	<b>41.1</b>	0.5	<b>9654</b>	<b>39.7</b>	0.5	<b>1407</b>	<b>61.8</b>	1.7	<b>46.8</b>	2.0	<b>7.8</b>	1.0	<b>45.4</b>	1.9	<b>0.2</b>	0.650
<b>Any traumatic events</b>	<b>18535</b>	<b>71.8</b>	0.4	<b>16513</b>	<b>70.5</b>	0.5	<b>2022</b>	<b>90.4</b>	1.1	<b>75.8</b>	1.5	<b>5.0</b>	0.7	<b>19.2</b>	1.5	<b>337.3*</b>	<.001

<sup>a</sup>Chi square tests comparing the proportion of respondents with TE onset prior to PE onset versus the proportion of respondents with PE onset prior to TE onset. \*Significant at the .05 level, 2-sided test.

<sup>b</sup>Estimates are based on weighted data.

<sup>c</sup>Brazil did not ask about this trauma.

<sup>d</sup>New Zealand did not ask about this trauma.

<sup>e</sup>Nigeria, Lebanon, Belgium, France, Germany, Italy, the Netherlands and Spain did not ask about this trauma.

<sup>f</sup>Goodness-of-fit test cannot be computed because of small expected counts (<5).



# BJPpsych

The British Journal of Psychiatry

## Trauma and psychotic experiences: transnational data from the World Mental Health Survey

John J. McGrath, Sukanta Saha, Carmen C. W. Lim, Sergio Aguilar-Gaxiola, Jordi Alonso, Laura H. Andrade, Evelyn J. Bromet, Ronny Bruffaerts, José M. Caldas de Almeida, Graça Cardoso, Giovanni de Girolamo, John Fayyad, Silvia Florescu, Oye Gureje, Josep M. Haro, Norito Kawakami, Karestan C. Koenen, Viviane Kovess-Masfety, Sing Lee, Jean-Pierre Lepine, Katie A. McLaughlin, Maria E. Medina-Mora, Fernando Navarro-Mateu, Akin Ojagbemi, Jose Posada-Villa, Nancy Sampson, Kate M. Scott, Hisateru Tachimori, Margreet ten Have, Kenneth S. Kendler, Ronald C. Kessler and on behalf of the WHO World Mental Health Survey Collaborators

BJP published online November 2, 2017 Access the most recent version at DOI:

[10.1192/bjp.bp.117.205955](https://doi.org/10.1192/bjp.bp.117.205955)

---

### Supplementary Material

Supplementary material can be found at:

<http://bjp.rcpsych.org/content/suppl/2017/10/31/bjp.bp.117.205955.DC1>

### References

This article cites 0 articles, 0 of which you can access for free at:

<http://bjp.rcpsych.org/content/early/2017/10/25/bjp.bp.117.205955#BIBL>

### Reprints/permissions

To obtain reprints or permission to reproduce material from this paper, please write to [permissions@rcpsych.ac.uk](mailto:permissions@rcpsych.ac.uk)

### P<P

Published online 2017-11-02T00:05:09-07:00 in advance of the print journal.

### You can respond to this article at

[/letters/submit/bjprcpsych;bjp.bp.117.205955v1](http://bjp.rcpsych.org/letters/submit/bjprcpsych;bjp.bp.117.205955v1)

### Downloaded from

<http://bjp.rcpsych.org/> on December 14, 2017  
Published by The Royal College of Psychiatrists

---

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

---

To subscribe to *The British Journal of Psychiatry* go to:  
<http://bjp.rcpsych.org/site/subscriptions/>

---